

Importance of neutrophil-to-lymphocyte ratio in coronary artery disease

Koroner arter hastalığında nötrofil-lenfosit oranının önemi

Kerem İnanoğlu

Department of Anesthesiology and Reanimation, Health Sciences University Antalya Training and Research Hospital, Antalya, Turkey

With great interest, I have read the article by Coskun et al (1).

The authors report that there is a correlation between mean platelet volume (MPV) and Gensini score; however, there is no significant correlation between neutrophil-to-lymphocyte (N/L) ratio and Gensini score in myocardial infarction associated with ST elevation. They also report that there is no significant correlation between both N/L ratio and MPV and Gensini score in myocardial infarction without ST elevation. However, studies in literature have shown the relationship between cardiovascular diseases and N/L ratio.

Leukocyte counts and the ratios of leukocyte subtypes are accepted as markers of inflammation in cardiovascular diseases. In relation to this, it has been shown that the N/L ratio is a marker of prognosis in heart failure, stable angina pectoris, and acute coronary syndromes. Elevated N/L ratios may reflect an inflammatory state and may be associated with perioperative myocardial damages and long term adverse outcomes. Neutrophils have been known to play a role in influencing the progression of atherosclerotic plaques. A reduced lymphocyte count is important to reflect physiological stress. A study reports the relationship between N/L ratios and mortality in ischemic heart disease (2). Similarly, increased preoperative N/L ratio is associated with increased mortality and morbidity as an independent risk factor in coronary artery bypass graft surgery (CABG) (3). Anesthesia methods used in patients undergoing CABG are also affected by N/L ratios (4).

Considering that increased preoperative N/L ratio is associated with long-term mortality after CABG (3) and preoperative N/L ratio is an independent predictor of saphenous vein graft patency after CABG (5), it becomes important to evaluate the preoperative N/L ratio, especially in patients undergoing CABG.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Çıkar Çatışması: Yazar çıkar çatışması bildirmemiştir.

Finansal Destek: Yazar bu çalışma için finansal destek almadıklarını beyan etmiştir.

REFERENCES

1. Coşkun FY, Sucu M, Aksoy N, Özer O. The neutrophile to lymphocyte ratio and mean platelet volume with Gensini score in patients with acute myocardial infarction. *Gaziantep Med J* 2015; 21: 200-4. [\[CrossRef\]](#)
2. Tamhane UU, Aneja S, Montgomery D, Rogers EK, Eagle KA, Gurm HS. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *Am J Cardiol* 2008; 102: 653-7. [\[CrossRef\]](#)
3. Gibson PH, Croal BL, Cuthbertson BH, Small GR, Ifezulike AI, Gibson G, et al. Preoperative neutrophil-lymphocyte ratio and outcome from coronary artery bypass grafting. *Am Heart J* 2007; 154: 995-1002. [\[CrossRef\]](#)
4. Aldemir M, Dogan BE, Adali F, Tecer E, Ozturk KN, Kavakli AS, et al. Comparison of the effects of propofol and desflurane anaesthesia on the neutrophil/lymphocyte ratio after coronary artery bypass surgery. *J Turgut Ozal Med Cent* 2015; 22: 165-70.
5. Taşoğlu I, Turak O, Nazli Y, Ozcan F, Colak N, Sahin S, et al. Preoperative neutrophil-lymphocyte ratio and saphenous vein graft patency after coronary artery bypass grafting. *Clin Appl Thromb Hemost* 2014; 20: 819-24. [\[CrossRef\]](#)

How to cite:

İnanoğlu K. Importance of neutrophil-to-lymphocyte ratio in coronary artery disease. *Eur J Ther* 2017; 23: 130.