

**Manuscript type:** Original Article

DOI: 10.5152/EurJTher.2019.600

**Title:** Congenital Extrahepatic Portosystemic Shunt: Abernethy Malformation Type 2

**Running head:** Abernethy Malformation Type 2

**Authors:** 1Ömer TOPRAK, 2Mehmet Nur KAYA, 2Sultan TÜREL, 2Uğur ERGÜN, 3Taha Yasin ARSLAN

**Instutions:** 1University of Balıkesir School of Medicine, Nephrology, Balıkesir, Turkey

2University of Balıkesir School of Medicine, Internal Medicine, Balıkesir, Turkey

3University of Balıkesir School of Medicine, Radiology, Balıkesir, Turkey

**Address for correspondence:** Mehmet Nur KAYA

**E-mail:** [mehmetnurkaya@yahoo.com](mailto:mehmetnurkaya@yahoo.com)

**Accepted:** 20.04.2018

**Cite this article as:** Toprak Ö, Kaya MN, Türel S, Ergün U, Arslan TY. Congenital Extrahepatic Portosystemic Shunt: Abernethy Malformation Type 2. Eur J Ther 2019; 10.5152/EurJTher.2019.600

## INTRODUCTION

Abernethy malformation (AM) with congenital anomaly was defined by John Abernethy in 1973. This malformation is characterized by shunting and is between the portal vein (PV) and the systemic circulation (1, 2). AM is frequently associated with other rare congenital anomalies, including the extrahepatic portocaval shunt, heterotaxy, biliary atresia and liver nodules (3,4). There are two types of AM which have been classified into two types based on the pattern of anastomosis between the systemic circulation and PV and the presence an intrahepatic portal venous supply. AM Type I portosystemic shunts is characterized by complete shunting and absence of a PV. AM type II is characterized by partial shunting with a small grade of PV flow to the liver (5). Assessment of the vascular anatomy and liver by new abdominal imaging technological in AM aids planning, so that the patient can receive appropriate treatment. Alternative treatment mode non-surgical endovascular treatment, if this treatment fails, the liver transplantation may be considered (5). In this study, we present the case of a male who was incidentally diagnosed with AM of type II.

## CASE PRESENTATION

A 68-year-old male patient presented to the gastroenterology department with complaints of abdominal pain and nausea. The patient had no significant personal or family medical history. No pathology was found except the right upper quadrant sensitivity on physical examination. The patient underwent total gastrectomy for gastric adenocarcinoma 3 months ago. Laboratory tests showed hemoglobin 13 g/dL, direct bilirubin 1.24  $\mu\text{mol/L}$ , indirect bilirubin 3.70  $\mu\text{mol/L}$ , lactate dehydrogenase 322 U/L, alanine aminotransferase 21 U/L, aspartate aminotransferase 40 U/L, gamma-glutamyl transpeptidase 32 U/L, alkaline phosphatase 116 U/L, prothrombin time 39.0 sec. Serologic tests for hepatitis B and C viruses negative results. Abdominal ultrasonography showed the presence of an anechoic tubular structure approximately 13 mm in diameter in the liver with no current Doppler signal. Computed tomography (CT) axial sections showed superior contrast enhancement in the right lobe of the liver, suggesting a 5 mm diameter hemangioma. PV diameter was measured as 5 mm. PV superior mesenteric vein (SMV) and splenic vein junction left renal vein portocaval shunt. (Figure A, B) Given these radiological findings, the hypoplastic PV with a portocaval shunt (AM of type II), The patient continued conservative treatment and after 1-month abdominal pain and nausea symptom was mild. At 3-month follow-up, he was no change on ultrasonography. Firstly the patient will consider the alternative treatment mode non-surgical endovascular treatment. If this treatment fails, the liver transplantation may be considered. (Informed consent is taken from the patient before writing this report)

## DISCUSSION

Congenital anomalies and vascular shunt diseases are seen together (6). AM can be anatomically classified with radiological imaging techniques. AM type II portosystemic shunt is characterized by the presence of a patent intrahepatic portal venous supply and a partial shunt (7). Type I can be further subclassified into type Ia and type Ib. Type Ia, separate drainage of the SMV and splenic vein into systemic veins, type Ib where SMV and splenic vein join to form a short extrahepatic portal vein which drains into a systemic vein. This patient had a type II AM with a side-to-side portocaval shunt between the left renal vein and splenic vein. Congenital vascular malformations are frequently associated with congenital anomalies. Other anomalies have also been reported in patients with AM which include chromosomal anomalies such as Down syndrome and structural anomalies of the cardiac defects, biliary atresia, polysplenia, situs inversus (8,9,10). Hepatic shunt frequently can also present with hypoglycemia. This might be the effect of defective uptake of glucose and defect of insulin secretion due to reduced hepatic degradation of the normal quantity of secreted insulin (11,12). A diagnosis of AM can now be made by noninvasive abdominal imaging technologies, such as USG, CT, and Magnetic Resonance Imaging (MRI) (13). The imaging findings in patients with AM with hepatocellular carcinoma do not appear to be typical, that is hypervascularity on the arterial phase images with washout on delayed phase (14). Patients who do not have typical findings of a benign lesion, i.e. lack arterial enhancement or arterial enhancement without washout, should be closely followed up. Two groups according to the type of shunt those should be offered shunt closure either interventional embolization or surgical whereas those with type 1 shunts should be liver transplanted (15,16).

## CONCLUSION

AM is a rare congenital vascular malformation that can be diagnosed to abdominal imaging technological (USG, CT, MRI). In this case, we present the case of a male who was incidentally diagnosed with AM of type II. Endovascular treatment of Abernethy malformation type 2 should be the first-line treatment if not successful liver transplantation should be considered

## REFERENCES

1. Howard ER, Davenport M. Congenital extrahepatic portocaval shunts the Abernethy malformation. *J Pediatr Surg* 1997; 32(3): 494–7.
2. Loomba RS, Telega GW, Gudausky TM. Type 2 Abernethy malformation presenting as a portal vein-coronary sinus fistula. *J Pediatr Surg* 2012; 47(5): E25–31.
3. Schaeffer DF, Laiq S, Jang HJ. Abernethy malformation type II with nephrotic syndrome and other multisystemic presentation: an illustrative case for understanding pathogenesis of extrahepatic complication of the congenital portosystemic shunt. *Hum Pathol* 2013; 44(3): 432–7.
4. Hao Y, Hong X, Zhao X. Congenital absence of the portal vein associated with focal nodular hyperplasia of the liver and congenital heart disease (Abernethy malformation): a case report and literature review. *Oncol Lett* 2015; 9(2): 695–700.
5. Kong Yalin, Zhang Hongyi, Liu Chengli. Abernethy malformation with multiple aneurysms: incidentally found in an adult woman with Caroli's disease. *Annals of Hepatology* Vol. 12 No.2, 2013: 327-331
6. Passalacqua M, Lie KT, Yarmohammadi H. Congenital extrahepatic portosystemic shunt (Abernethy malformation) treated endovascularly with vascular plug shunt closure. *Pediatric Surg International* 2012; 1: 79-83.
7. Mistinova J, Valassis F, Varga I. Congenital absence of the portal vein-Case report and a review of the literature. *Clin Anat* 2010; 7: 750-8.
8. Marchetti F, Pellegrino L, Al-Ansari N. Congenital absence of the portal vein in a middle-aged man. *Surg Radiol Anat* 2011; 33: 369-72.
9. Hu GH, Shen LG, Yang J. Insight into the congenital absence of the portal vein: Is it rare? *World J Gastroenterol* 2008; 14: 5969-79.
10. Franchi-Abella S, Branchereau S, Lambert V. Complications of congenital portosystemic shunts in children: Therapeutic options and outcomes. *J Pediatr Gastroenterol Nutr* 2010; 51: 322-30

11. Ratnasamy C, Kurbegov A, Swaminathan S. Cardiac anomalies in the setting of the Abernethy malformation of the portal vein. *Cardiol Young* 2007; 17(2): 212–214,
12. Shinkai M, Ohhama Y, Nishi T. Congenital absence of the portal vein and role of liver transplantation in children. *J Pediatr Surg* 2001; 36(7): 1026–1031
13. Konstas AA, Digumarthy SR, Avery LL. Congenital portosystemic shunts: Imaging findings and clinical presentations in 11 patients. *Eur J Radiol* 2011; 2: 175-81.
14. Morgan G, Superina R. Congenital absence of the portal vein: Two cases and a proposed classification system for portosystemic vascular anomalies. *J Pediatr Surg* 1994; 29: 1239-41.
15. Gallego C, Miralles M, Marín C. Congenital hepatic shunts. *Radiographics* 2004; 24: 755-72.
16. Akahoshi T, Nishizaki T, Wakasugi K. Portal-systemic encephalopathy due to a congenital extrahepatic portosystemic shunt: Three cases and literature review. *Hepatogastroenterology* 2000; 47: 1113-6.

**Figure A-B.** CT axial sections showed the SMV and the splenic vein merges into the left renal vein drain. PV is thin-walled and hypoplastic.



UNCONFIRMED

